

Discrete Comput Geom (2015) 53:221–225
DOI 10.1007/s00454-014-9645-y

Triangle-Free Geometric Intersection Graphs with No Large Independent Sets

Bartosz Walczak

Received: 1 April 2014 / Revised: 12 October 2014 / Accepted: 27 October 2014 /

Published online: 12 November 2014

© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract It is proved that there are triangle-free intersection graphs of line segments in the plane with arbitrarily small ratio between the maximum size of an independent set and the total number of vertices.

Keywords Intersection graph · Line segments · Triangle-free · Maximum independent set · Fractional chromatic number

1 Introduction

Pawlik et al. [7] proved that there are triangle-free intersection graphs of line segments in the plane with arbitrarily large chromatic number. The graphs they construct have independent sets containing more than $1/3$ of all the vertices. It has been left open whether there is a constant $c > 0$ such that every triangle-free intersection graph of n segments in the plane has an independent set of size at least cn . Fox and Pach [3] conjectured a much more general statement, that K_k -free intersection graphs of curves in the plane have linear-size independent sets, for every k . This would imply a well-known conjecture that k -quasi-planar graphs (graphs drawn in the plane so that no k edges cross each other) have linearly many edges [5], which is proved up to $k = 4$ [1].

In this note, I resolve the independent set problem in the negative, proving the following strengthening of the result of Pawlik et al.:

Theorem *There are triangle-free segment intersection graphs with arbitrarily small ratio between the maximum size of an independent set and the total number of vertices.*

B. Walczak

Theoretical Computer Science Department, Faculty of Mathematics and Computer Science,
Jagiellonian University, Kraków, Poland
e-mail: walczak@tcs.uj.edu.pl

The constructions presented in the next two sections give rise to triangle-free intersection graphs of n segments in the plane with maximum independent set size $\Theta(n/\log \log n)$.

2 Construction

Pawlik et al. [7] construct, for $k \geq 1$, a triangle-free graph G_k and a family \mathcal{P}_k of subsets of $V(G_k)$, called *probes*, with the following properties:

- (i) $|\mathcal{P}_k| = 2^{2^{k-1}-1}$,
- (ii) every member of \mathcal{P}_k is an independent set of G_k ,
- (iii) for every proper coloring of the vertices of G_k , there is a probe $P \in \mathcal{P}_k$ such that at least k colors are used on the vertices in P .

They are built by induction on k , as follows. The graph G_1 has just one vertex v , and \mathcal{P}_1 has just one probe $\{v\}$. For $k \geq 2$, first, a copy (G, \mathcal{P}) of $(G_{k-1}, \mathcal{P}_{k-1})$ is taken. Then, for every probe $P \in \mathcal{P}$, another copy (G_P, \mathcal{P}_P) of $(G_{k-1}, \mathcal{P}_{k-1})$ is taken. There are no edges between vertices from different copies. Finally, for every probe $P \in \mathcal{P}$ and every probe $Q \in \mathcal{P}_P$, a new vertex d_Q connected to all vertices in Q , called the *diagonal* of Q , is added. The resulting graph is G_k . The family of probes \mathcal{P}_k is defined by

$$\mathcal{P}_k = \{P \cup Q : P \in \mathcal{P} \text{ and } Q \in \mathcal{P}_P\} \cup \{P \cup \{d_Q\} : P \in \mathcal{P} \text{ and } Q \in \mathcal{P}_P\}.$$

It is easy to check that the graph G_k is indeed triangle-free and the conditions (i)–(iii) are satisfied for (G_k, \mathcal{P}_k) —see [7] for details. It is also shown in [7] how the graph G_k is represented as a segment intersection graph.

I will show that there is an assignment w_k of positive integer weights to the vertices of G_k with the following properties:

- (iv) the total weight of G_k is $\frac{k+1}{2} \cdot 2^{2^{k-1}-1}$,
- (v) for every independent set I of G_k , the number of probes $P \in \mathcal{P}_k$ such that $P \cap I \neq \emptyset$ is at least the weight of I .

Once this is achieved, the proof of the theorem of this paper follows easily. Namely, it follows from (i) and (v) that every independent set I of G_k has weight at most $2^{2^{k-1}-1}$. We can take the representation of G_k as a segment intersection graph and replace every segment representing a vertex $v \in V(G_k)$ by $w_k(v)$ parallel segments lying very close to each other, so as to keep the property that any two segments representing vertices $u, v \in V(G_k)$ intersect if and only if $uv \in E(G_k)$. It follows from (iv) that the family of segments obtained this way has size $\frac{k+1}{2} \cdot 2^{2^{k-1}-1}$, while every independent set of its intersection graph has size at most $2^{2^{k-1}-1}$.

The assignment w_k of weights to the vertices of G_k is defined by induction on k , following the inductive construction of (G_k, \mathcal{P}_k) . The weight of the only vertex of G_1 is set to 1. This clearly satisfies (iv) and (v). For $k \geq 2$, let $G, \mathcal{P}, G_P, \mathcal{P}_P$ and d_Q be defined as in the inductive step of the construction of (G_k, \mathcal{P}_k) . Let $p = |\mathcal{P}_{k-1}| = 2^{2^{k-2}-1}$. The weights w_k of the vertices of G are their original weights w_{k-1} in G_{k-1} multiplied by p . The weights w_k of the vertices of every G_P are equal to their original

weights w_{k-1} in G_{k-1} . The weight w_k of every diagonal d_Q is set to 1. It remains to prove that (iv) and (v) are satisfied for $(G_k, \mathcal{P}_k, w_k)$ assuming that they hold for $(G_{k-1}, \mathcal{P}_{k-1}, w_{k-1})$.

The proof of (iv) is straightforward:

$$w_k(G_k) = w_k(G) + \sum_{P \in \mathcal{P}} (w_k(G_P) + |\mathcal{P}_P|) = 2pw_{k-1}(G_{k-1}) + p^2 = \frac{k+1}{2} \cdot 2^{2^{k-1}-1}.$$

For the proof of (v), let I be an independent set in G_k . Let $\mathcal{I} = \{P \in \mathcal{P} : P \cap I \neq \emptyset\}$. For every probe $P \in \mathcal{P}$, define

$$\begin{aligned} \mathcal{I}_P &= \{Q \in \mathcal{P}_P : Q \cap I \neq \emptyset\}, & \mathcal{P}'_P &= \{P \cup Q : Q \in \mathcal{P}_P\} \cup \{P \cup \{d_Q\} : Q \in \mathcal{P}_P\}, \\ D_P &= \{d_Q : Q \in \mathcal{P}_P\}, & \mathcal{I}'_P &= \{P' \in \mathcal{P}'_P : P' \cap I \neq \emptyset\}. \end{aligned}$$

By the induction hypothesis, we have

$$w_k(V(G) \cap I) \leq p|\mathcal{I}|, \quad w_k(V(G_P) \cap I) \leq |\mathcal{I}_P|.$$

Suppose $P \in \mathcal{I}$. It follows that $(P \cup Q) \cap I \neq \emptyset$ and $(P \cup \{d_Q\}) \cap I \neq \emptyset$ for every $Q \in \mathcal{P}_P$. Hence $|\mathcal{I}'_P| = |\mathcal{P}'_P| = 2p$. Moreover, we have $d_Q \notin I$ whenever $Q \in \mathcal{I}_P$, because d_Q is connected to all vertices in Q , one of which belongs to I . Hence

$$w_k(V(G_P) \cap I) + w_k(D_P \cap I) \leq |\mathcal{I}_P| + |\mathcal{P}_P \setminus \mathcal{I}_P| = |\mathcal{P}_P| = p.$$

Now, suppose $P \in \mathcal{P} \setminus \mathcal{I}$. If $Q \in \mathcal{I}_P$, then $(P \cup Q) \cap I \neq \emptyset$, $d_Q \notin I$ (by the same argument as above), and $(P \cup \{d_Q\}) \cap I = \emptyset$. If $Q \in \mathcal{P}_P \setminus \mathcal{I}_P$, then $(P \cup Q) \cap I = \emptyset$, and $(P \cup \{d_Q\}) \cap I \neq \emptyset$ if and only if $d_Q \in I$. Hence

$$w_k(V(G_P) \cap I) + w_k(D_P \cap I) \leq |\mathcal{I}_P| + |D_P \cap I| = |\mathcal{I}'_P|.$$

To conclude, we gather all the inequalities and obtain

$$\begin{aligned} w_k(I) &= w_k(V(G) \cap I) + \sum_{P \in \mathcal{P}} (w_k(V(G_P) \cap I) + w_k(D_P \cap I)) \\ &\leq p|\mathcal{I}| + \sum_{P \in \mathcal{I}} p + \sum_{P \in \mathcal{P} \setminus \mathcal{I}} |\mathcal{I}'_P| = \sum_{P \in \mathcal{I}} |\mathcal{I}'_P| + \sum_{P \in \mathcal{P} \setminus \mathcal{I}} |\mathcal{I}'_P| = \sum_{P \in \mathcal{P}} |\mathcal{I}'_P|. \end{aligned}$$

3 Improved Construction

Pawlik et al. [7] define also a graph \tilde{G}_k , which arises from (G_k, \mathcal{P}_k) by adding, for every probe $P \in \mathcal{P}_k$, a diagonal d_P connected to all vertices in P . This is the smallest triangle-free segment intersection graph known to have chromatic number greater than k . Define the assignment \tilde{w}_k of weights to the vertices of \tilde{G}_k so that \tilde{w}_k is equal to w_k on the vertices of G_k and $\tilde{w}_k(d_P) = 1$ for every $P \in \mathcal{P}_k$. Let I be an independent set in \tilde{G}_k . Let $\mathcal{I} = \{P \in \mathcal{P}_k : P \cap I \neq \emptyset\}$. Hence $d_P \notin I$ for $P \in \mathcal{I}$. It follows that

$$\begin{aligned}
\tilde{w}_k(I) &= w_k(V(G_k) \cap I) + |\{d_P : P \in \mathcal{P}_k\} \cap I| \\
&\leq |\mathcal{I}| + |\mathcal{P}_k \setminus \mathcal{I}| = |\mathcal{P}_k| = 2^{2^{k-1}-1}, \\
\tilde{w}_k(\tilde{G}_k) &= w_k(G_k) + |\mathcal{P}_k| = \frac{k+3}{2} \cdot 2^{2^{k-1}-1}.
\end{aligned}$$

The graph \tilde{G}_k is the smallest one for which I can prove that it has a weight assignment such that the ratio between the maximum weight of an independent set and the total weight is at most $\frac{2}{k+3}$. It is not difficult to prove (e.g. using weak LP duality) that the assignment of weights \tilde{w}_k to the vertices of \tilde{G}_k is optimal (gives the least ratio) for this particular graph.

Both constructions give rise to triangle-free intersection graphs of n segments in the plane with maximum independent set size $\Theta(n/\log \log n)$. On the other hand, it follows from the result of McGuinness [4] that every triangle-free intersection graph of n segments has chromatic number $O(\log n)$ and maximum independent set size $\Omega(n/\log n)$.

4 Other Geometric Shapes

It is known that the graphs G_k and \tilde{G}_k have intersection models by many other geometric shapes, for example, L-shapes, axis-parallel ellipses, circles, axis-parallel square boundaries [6] or axis-parallel boxes in \mathbb{R}^3 [2]. The result of this paper can be extended to those models for which every geometric object X representing a vertex of the intersection graph can be replaced by many pairwise disjoint objects approximating X . This is possible, for example, for intersection graphs of L-shapes, circles or axis-parallel square boundaries, but not for intersection graphs of axis-parallel ellipses or axis-parallel boxes in \mathbb{R}^3 . The problem whether triangle-free intersection graphs of the latter kind of shapes have linear-size independent sets remains open.

Acknowledgments The author thanks Michael Hoffmann for helpful discussions. The work was supported by Polish National Science Center Grant 2011/03/B/ST6/01367 and Swiss National Science Foundation Grant 200020-144531.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

1. Ackerman, E.: On the maximum number of edges in topological graphs with no four pairwise crossing edges. *Discrete Comput. Geom.* **41**(3), 365–375 (2009)
2. Burling, J. P.: On coloring problems of families of prototypes. Ph.D. Thesis, University of Colorado, Boulder (1965)
3. Fox, J., Pach, J.: Coloring K_k -free intersection graphs of geometric objects in the plane. *Eur. J. Comb.* **33**(5), 853–866 (2012)
4. McGuinness, S.: Colouring arcwise connected sets in the plane I. *Graphs Comb.* **16**(4), 429–439 (2000)
5. Pach, J., Shahrokhi, F., Szegedy, M.: Applications of the crossing number. *Algorithmica* **16**(1), 111–117 (1996)

6. Pawlik, A., Kozik, J., Krawczyk, T., Lasoń, M., Micek, P., Trotter, W.T., Walczak, B.: Triangle-free geometric intersection graphs with large chromatic number. *Discrete Comput. Geom.* **50**(3), 714–726 (2013)
7. Pawlik, A., Kozik, J., Krawczyk, T., Lasoń, M., Micek, P., Trotter, W.T., Walczak, B.: Triangle-free intersection graphs of line segments with large chromatic number. *J. Comb. Theory, Ser. B* **105**, 6–10 (2014)